

Nr. 95  
10. May 2023

Preprint-Series: Department of Mathematics - Applied Mathematics

Translation invariant diagonal frame decomposition  
for the Radon transform

S. Göppel, J. Friel, M. Haltmeier

AppliedMathematics

---

Technikerstraße 13 - 6020 Innsbruck - Austria  
Tel.: +43 512 507 53803 Fax: +43 512 507 53898  
<https://applied-math.uibk.ac.at>

# Translation invariant diagonal frame decomposition for the Radon transform

Simon Göppel

Department of Mathematics, University of Innsbruck  
Technikerstrasse 13, 6020 Innsbruck, Austria  
E-mail: [simon.goeppel@uibk.ac.at](mailto:simon.goeppel@uibk.ac.at)

Jürgen Friel

Department of Computer Science and Mathematics, OTH Regensburg  
Galgenbergstraße 32, 93053 Regensburg, Germany  
E-mail: [juergen.friel@oth-regensburg.de](mailto:juergen.friel@oth-regensburg.de)

Markus Haltmeier

Department of Mathematics, University of Innsbruck  
Technikerstrasse 13, 6020 Innsbruck, Austria  
E-mail: [markus.haltmeier@uibk.ac.at](mailto:markus.haltmeier@uibk.ac.at)

May 10, 2023

## Abstract

In this article, we address the challenge of solving the ill-posed reconstruction problem in computed tomography using a translation invariant diagonal frame decomposition (TI-DFD). First, we review the concept of a TI-DFD for general linear operators and the corresponding filter-based regularization concept. We then introduce the TI-DFD for the Radon transform on  $L^2(\mathbb{R}^2)$  and provide an exemplary construction using the TI wavelet transform. Presented numerical results clearly demonstrate the benefits of our approach over non-translation invariant counterparts.

**Keywords:** DFD, translation invariance, Radon transform, inverse problems, regularization

# 1 Introduction

Computed tomography (CT) is a widely applied imaging modality in medicine and industry, where the underlying mathematical model is the Radon transform. For a function or signal  $f: \mathbb{R}^2 \rightarrow \mathbb{R}$ , the Radon transform is denoted by  $\mathcal{R}f: \mathbb{S}^1 \times \mathbb{R} \rightarrow \mathbb{R}$ . It is well-known, that reconstructing a function  $f$  from approximate knowledge of its line integrals amounts to an ill-posed inverse problem of the form  $g^\delta = \mathcal{R}f + \eta$  [13]. Here,  $\delta > 0$  is some noise level and  $\eta$  denotes the data distortions with  $\|\eta\| \leq \delta$ . In particular, this means that inverting the Radon transform is unstable and that small perturbations in data can cause big reconstruction errors.

Classical filter based regularization is a well-known stabilization concept. Assuming a singular value decomposition (SVD)  $\mathcal{R}f = \sum_{n \in \mathbb{N}} \sigma_n \langle f, u_n \rangle v_n$  for the Radon transform, a regularized reconstruction is given by  $f_\alpha^\delta = \sum_{n \in \mathbb{N}} \Phi_\alpha(\sigma_n) \langle g^\delta, v_n \rangle u_n$ , depending on a so-called regularizing filter  $\Phi_\alpha$ , see [5, 8]. However, filtering based on the SVD comes with several shortcomings. In general, for an arbitrary linear operator, the SVD might be hard to compute numerically or not be known analytically. Additionally, the basis elements  $u_n, v_n$  are only adapted to the operator itself, but not the underlying signal class of interest.

To overcome these limitations, the so-called diagonal frame decomposition (DFD) as a generalization of the SVD has been studied thoroughly in the recent years. In particular, DFDs are better suited as they not only can be adjusted to the underlying application and involved signals, but often also provide efficient implementations. A prominent example of a DFD for the Radon transform is the wavelet-vaguelette decomposition (WVD) introduced in [4]. Related construction involving curvelets and shearlets can for example be found in [1, 3]. A more general analysis of regularization properties and convergence results have been presented in [5, 6, 9].

One drawback of the classical WVD reconstruction approach is that in general it lacks translation invariance, which can lead to well-known wavelet artifacts in the reconstruction [11]. Translation invariant systems on the other hand are known to perform better in that regard for simple tasks such as denoising [2, 12]. To overcome this, in [7] the authors have introduced the concept of the translation invariant diagonal frame decomposition (TI-DFD) for general linear operators. Along with an analysis of the regularization properties of the filter based TI-DFD, the authors gave an exemplary construction of a TI-WVD regarding stable differentiation. These findings indicate improved regularization properties, when translation invariance is restored in the underlying wavelet system. Thus, the goal of the present paper is to construct a TI-WVD for the Radon transform. This way, we obtain an explicit filter based regularization strategy, which can be implemented efficiently. We will demonstrate its improved regularization properties by comparing the results to the classical WVD in a numerical example.

**Notation.** For  $f \in L^2(\mathbb{R}^2)$ , the Fourier transform is denoted by  $\widehat{f} = \mathbf{F}f$ , where  $\widehat{f}(\xi) := \int_{\mathbb{R}^2} f(x)e^{-i\langle \xi, x \rangle}$  if  $f \in L^2(\mathbb{R}^d) \cap L^1(\mathbb{R}^d)$ . For  $f \in L^2(\mathbb{S} \times \mathbb{R})$ , the one dimensional Fourier transform in the second variable is denoted by  $\mathbf{F}_2$ . Furthermore, we write  $u^*(x) := \overline{u(-x)}$ , where  $\bar{z}$  is the complex conjugate of  $z \in \mathbb{C}$ .

## 2 The Translation Invariant Diagonal Frame Decomposition

In this section, we recall the concept of translation invariant (TI) frames [11] and translation invariant diagonal frame decompositions (TI-DFDs) of linear operators [7]. Furthermore, we recall the concept of filtered regularization using TI-DFDs.

**Definition 2.1** (TI-frame). *Let  $\Lambda$  be an at most countable index set. We call the family  $(u_\lambda)_{\lambda \in \Lambda} \in L^2(\mathbb{R}^d)^\Lambda$  a translation invariant frame (TI-frame) for  $L^2(\mathbb{R}^d)$  if for all  $\lambda \in \Lambda$  we have  $\widehat{u}_\lambda \in L^\infty(\mathbb{R}^d)$  and there exist constants  $A, B > 0$ , such that*

$$\forall f \in L^2(\mathbb{R}^d): \quad A\|f\|^2 \leq \sum_{\lambda \in \Lambda} \|u_\lambda^* * f\|^2 \leq B\|f\|^2. \quad (2.1)$$

We call a TI-frame  $(u_\lambda)_{\lambda \in \Lambda}$  tight if (2.1) holds with TI-frame bounds  $A = B = 1$ .

Defining  $w_\lambda := \mathbf{F}^{-1}(2\pi\widehat{u}_\lambda / \sum_{\mu \in \Lambda} |\widehat{u}_\mu|^2)$  for every TI frame element  $u_\lambda$ , we obtain the so-called canonical dual  $(w_\lambda)_{\lambda \in \Lambda}$  of  $(u_\lambda)_{\lambda \in \Lambda}$ . It holds that

$$\forall f \in L^2(\mathbb{R}^d): \quad f = \sum_{\lambda \in \Lambda} w_\lambda * (u_\lambda^* * f). \quad (2.2)$$

Note that (2.2) in fact holds true for any dual frame  $(w_\lambda)_{\lambda \in \Lambda}$  defined by the property  $\sum_{\lambda} (\mathbf{F}w_\lambda) \cdot (\overline{\mathbf{F}u_\lambda}) = 2\pi$ . In particular, the canonical dual always exists but it not uniquely defined by (2.2). Further note that TI-frame is not a frame in the classical sense. In particular, TI frame coefficients  $u_\lambda^* * f(x) = \langle f, u_\lambda(\cdot - x) \rangle$  use a continuous translation parameter  $x$ .

**Definition 2.2** (TI-DFD). *Let  $\mathbf{K}: \mathcal{D}(\mathbf{K}) \subseteq L^2(\mathbb{R}^d) \rightarrow \mathbb{Y}$  be a closed linear operator and let the space of bounded operators between  $\mathbb{Y}$  and  $L^2(\mathbb{R}^d)$  be denoted by  $B(\mathbb{Y}, L^2(\mathbb{R}^d))$ . We call the system  $(u_\lambda, \mathcal{V}_\lambda^*, \kappa_\lambda)_{\lambda \in \Lambda}$  a translation invariant frame decomposition (TI-DFD) for  $\mathbf{K}$ , if the following hold:*

(TI1)  $(u_\lambda)_{\lambda \in \Lambda} \in L^2(\mathbb{R}^d)^\Lambda$  is a TI-frame for  $L^2(\mathbb{R}^d)$ .

(TI2)  $\forall \lambda \in \Lambda$  we have  $\mathcal{V}_\lambda^* \in B(\mathbb{Y}, L^2(\mathbb{R}^d))$  and

$$\forall g \in \overline{\text{ran } \mathbf{K}}: \sum_{\lambda \in \Lambda} \|\mathcal{V}_\lambda^* g\|^2 \asymp \|g\|^2.$$

(TI3)  $\forall \lambda \in \Lambda: \kappa_\lambda \in (0, \infty)$  and

$$\forall f \in \mathcal{D}(\mathbf{K}): \mathcal{V}_\lambda^*(\mathbf{K}f) = \kappa_\lambda (u_\lambda^* * f).$$

Here, we define  $F \asymp G \Leftrightarrow \exists c_1, c_2 > 0: c_1 G \leq F \leq c_2 G$ .

**Definition 2.3** (Regularizing filter). *A family  $(\Phi_\alpha)_{\alpha>0}$  of piecewise continuous functions  $\Phi_\alpha: (0, \infty) \rightarrow \mathbb{R}$  is called a regularizing filter if the following hold:*

(F1)  $\forall \alpha > 0: \|\Phi_\alpha\|_\infty < \infty$ .

(F2)  $\exists C > 0: \sup\{|\kappa \Phi_\alpha(\kappa)|: \alpha > 0 \wedge \kappa \geq 0\} \leq C$ .

(F3)  $\forall \kappa \in (0, \infty): \lim_{\alpha \rightarrow 0} \Phi_\alpha(\kappa) = 1/\kappa$ .

The following theorem summarizes main results of [7, Sections 2 and 3]. For that recall the notion of a regularization method [8, Definition 3.1].

**Theorem 2.4.** *Let  $(u_\lambda, \mathcal{V}_\lambda^*, \kappa_\lambda)_{\lambda \in \Lambda}$  be a TI-DFD for  $\mathbf{K}$ , let  $(w_\lambda)_{\lambda \in \Lambda}$  be a dual TI-frame for  $(u_\lambda)_{\lambda \in \Lambda}$  and*

$$\mathcal{R}_\alpha^\Phi g := \sum_{\lambda \in \Lambda} w_\lambda * (\Phi_\alpha(\kappa_\lambda) \cdot (\mathcal{V}_\lambda^* g)) \tag{2.3}$$

where  $(\Phi_\alpha)_{\alpha>0}$  is a regularizing filter. Then we have:

1.  $\forall g \in \text{ran}(\mathbf{K}): \mathbf{K}^{-1}g = \sum_{\lambda \in \Lambda} w_\lambda * (\kappa_\lambda^{-1} \cdot (\mathcal{V}_\lambda^* g))$ .

2. The family  $(\mathcal{R}_\alpha^\Phi)_{\alpha>0}$  together with suitable parameter choice, defines a regularization method for inverting  $\mathbf{K}$ .

The instability of inverting  $\mathbf{K}$  is reflected via the quasi-singular values  $(\kappa_\lambda)_{\lambda \in \Lambda}$ . More precisely, in [7] it has been shown that the inverse operator  $\mathbf{K}^{-1}$  is unbounded if and only if the quasi-singular values  $\kappa_\lambda$  accumulate at zero, assuming  $\inf_{\lambda} \|\mathcal{V}_\lambda^*\| > 0$ .

### 3 TI-DFD for the Radon Transform

Recall that for  $f \in L^1(\mathbb{R}^2) \cap L^2(\mathbb{R}^2)$  the Radon transform  $\mathcal{R}f: \mathbb{S} \times \mathbb{R} \rightarrow \mathbb{R}$  is defined by

$$\mathcal{R}f(\theta, s) = \int_{\mathbb{R}} f(s\theta + t\theta^\perp) dt, \quad (3.1)$$

for almost every  $(\theta, s) \in \mathbb{S}^1 \times \mathbb{R}$ . In this section we extend the Radon transform to a closed operator between  $L^2$  spaces [14] and then construct corresponding TI-DFDs.

#### 3.1 The Radon transform on $L^2$

In what follows, we will make extensive use of the Fourier Slice theorem which states that for all  $f \in L^1(\mathbb{R}^2) \cap L^2(\mathbb{R}^2)$  and almost every  $(\theta, \sigma) \in \mathbb{S}^1 \times \mathbb{R}$  we have

$$\mathbf{F}_2 \mathcal{R}f(\theta, \sigma) = \mathbf{F}f(\sigma\theta). \quad (3.2)$$

In fact we use an extension of (3.2) to the natural domain of definition  $\mathcal{D}(\mathcal{R}) \supseteq L^1(\mathbb{R}^2) \cap L^2(\mathbb{R}^2)$ . To this end we define the operator  $\mathcal{B}: \mathcal{D}(\mathcal{B}) \subseteq L^2(\mathbb{R}^2) \rightarrow L^2(\mathbb{S}^1 \times \mathbb{R})$  by

$$\forall (\theta, \sigma) \in \mathbb{S}^1 \times \mathbb{R}: \quad \mathcal{B}f(\theta, \sigma) := f(\sigma\theta) \quad (3.3)$$

According to (3.2),  $\mathcal{B}$  is the Fourier representation of the Radon transform on  $L^1(\mathbb{R}^2) \cap L^2(\mathbb{R}^2)$ . We next state some properties of  $\mathcal{B}$ , without proofs.

**Proposition 3.1** (Properties of  $\mathcal{B}$ ). *The operator  $\mathcal{B}$  as defined above satisfies the following:*

- (a)  $\mathcal{D}(\mathcal{B}) = \{f: \|\cdot\|^{-1/2}f \in L^2(\mathbb{R}^2)\}$
- (b)  $\mathcal{D}(\mathcal{B}^*) = \{g: |\cdot|^{-1/2}g(\theta, \cdot) \in L^2(\mathbb{S}^1 \times \mathbb{R})\}$
- (c)  $\mathcal{D}(\mathcal{B})$  and  $\mathcal{D}(\mathcal{B}^*)$  are dense in  $L^2$ , respectively.
- (d)  $\mathcal{B}$  is well-defined, linear, injective and unbounded.
- (e)  $\mathcal{B}^*g(\xi) = \|\xi\|^{-1}g(\xi/\|\xi\|, \|\xi\|)$ .

**Definition 3.2** (Radon transform on  $L^2$ ). *The operator  $\mathcal{R}: \mathcal{D}(\mathcal{R}) \subseteq L^2(\mathbb{R}^2) \rightarrow L^2(\mathbb{S}^1 \times \mathbb{R})$  defined as composition*

$$\mathcal{R}f := (\mathbf{F}_2^{-1} \circ \mathcal{B} \circ \mathbf{F})(f) \quad (3.4)$$

is called the Radon transform on  $L^2$ .

**Proposition 3.3** (Properties of  $\mathcal{R}$ ). *The Radon transform  $\mathcal{R}$  satisfies the following properties:*

- (a)  $\mathcal{R}$  is well-defined, linear, injective and unbounded.
- (b)  $\mathcal{R}$  is the closed extension of the operator defined by (3.1).
- (c)  $\mathcal{D}(\mathcal{R}) = \{f \in L^2(\mathbb{R}^2) \mid \mathbf{F}f \in \mathcal{D}(\mathcal{B})\}$ .
- (d)  $\mathcal{D}(\mathcal{R}^*) = \{g \in L^2(\mathbb{S}^1 \times \mathbb{R}) \mid \mathbf{F}_2g \in \mathcal{D}(\mathcal{B}^*)\}$ .
- (e)  $\mathcal{D}(\mathcal{R})$  and  $\mathcal{D}(\mathcal{R}^*)$  are dense in  $L^2$ , respectively.
- (f)  $\mathcal{R}^* = \mathbf{F}^{-1} \circ \mathcal{B}^* \circ \mathbf{F}_2$ .

Proposition 3.3 in particular states that the Radon transform  $\mathcal{R}$  is an unbounded operator. On the other hand, the restriction of  $\mathcal{R}$  to various closed subspaces is bounded. The most common case is the restriction of  $\mathcal{R}$  to  $L^2_D(\mathbb{R}^2) := \{f \in L^2(\mathbb{R}^2) : \text{supp}(f) \subseteq D\}$ . However, the restriction to functions with compact support poses a-priori assumptions that are not translation invariant. Further, with the above considerations we obtain the classical filtered backprojection formula (FBP) [13] between  $L^2$ -spaces, i.e.

$$\mathcal{R}^{-1}g = \frac{1}{4\pi}(\mathcal{R}^* \circ \mathcal{I}_1)(g), \quad (3.5)$$

where  $g = \mathcal{R}f$  and  $f \in \mathcal{D}(\mathcal{R})$  [14, 15]. Here, the operator  $\mathcal{I}_1f := \mathbf{F}^{-1}(|\cdot|\mathbf{F}f)$  is known as Riesz potential.

### 3.2 Necessary conditions

Before constructing a TI-DFD for the Radon transform on  $L^2(\mathbb{R}^2)$  we derive some necessary conditions.

**Lemma 3.4** (Necessary Conditions). *Let  $u_\lambda \in L^2(\mathbb{R}^2)$ ,  $v_\lambda \in L^2(\mathbb{S}^1 \times \mathbb{R})$  and  $\kappa_\lambda > 0$  satisfy  $\mathcal{R}^*v_\lambda^* = \kappa_\lambda u_\lambda^*$  and  $\hat{v}_\lambda, |\cdot|^{-1}\hat{v}_\lambda \in L^\infty(\mathbb{S}^1 \times \mathbb{R})$ , and define  $\mathcal{V}_\lambda^*g := \mathcal{R}^*(v_\lambda^* * g)$ .*

1.  $(\mathcal{V}_\lambda)_{\lambda \in \Lambda}$  satisfies (TI3).
2. If  $(u_\lambda, \mathcal{V}_\lambda^*, \kappa_\lambda)_{\lambda \in \Lambda}$  is a TI-DFD for  $\mathcal{R}$ , then

$$\forall \lambda \in \Lambda: \quad \hat{v}_\lambda(\theta, \sigma) = \kappa_\lambda |\sigma| \hat{u}_\lambda(\sigma\theta). \quad (3.6)$$

*Proof.* 1) Let  $f \in \mathcal{D}(\mathcal{R})$ . Then

$$\begin{aligned}
\kappa_\lambda(u_\lambda^* * f) &= \mathbf{F}^{-1}(\kappa_\lambda \widehat{u}_\lambda^* \cdot \widehat{f}) \\
&= \mathbf{F}^{-1}(\mathcal{B}^* \widehat{v}_\lambda^* \cdot \|\cdot\| \mathcal{B}^* \mathcal{B} \widehat{f}) \\
&= \mathbf{F}^{-1} \mathcal{B}^* (\widehat{v}_\lambda^* \cdot \widehat{\mathcal{R}f}) \\
&= \mathbf{F}^{-1} \mathcal{B}^* \mathbf{F}_2 (v_\lambda^* *_s \mathcal{R}f) \\
&= \mathcal{R}^* (v_\lambda^* *_s \mathcal{R}f) \\
&= \mathcal{V}_\lambda^* (\mathcal{R}f).
\end{aligned}$$

2) According to 1) and the definition of  $\mathcal{V}_\lambda^*$  we have  $\mathcal{R}^* (v_\lambda^* *_s \mathcal{R}f) = \kappa_\lambda (u_\lambda^* * f)$ . By applying the Fourier transform on both sides and since  $\mathcal{D}(\mathcal{R})$  is dense, we obtain (3.6).  $\square$

### 3.3 Construction of the TI-DFD

Now, let  $(u_\lambda)_{\lambda \in \Lambda}$  be a 2D (tensor product) TI wavelet frame with mother wavelet  $u \in L^2(\mathbb{R}^2)$ . That is, we assume the multi-scale structure

$$\forall (j, l) \in \Lambda := \mathbb{Z} \times \{\text{H}, \text{V}, \text{D}\}: u_{j,l}(x) = 2^j u_{0,l}(2^j x), \quad (3.7)$$

where  $j \in \mathbb{Z}$  is the scale index and  $l \in \{\text{H}, \text{V}, \text{D}\}$  indicates the horizontal, vertical or diagonal mother wavelet, respectively.

**Theorem 3.5** (TI-WVD for  $\mathcal{R}$ ). *Let  $(u_{j,l})_{j,l \in \Lambda}$  be defined by (3.7), suppose  $\text{supp}(\widehat{u}_{0,l}) = \{\xi \mid a \leq \|\xi\| \leq b\}$  for some  $a, b > 0$ , and for  $(j, l) \in \mathbb{Z} \times \{\text{H}, \text{V}, \text{D}\}$  define*

$$v_{j,l} := 2^{-j/2} \mathcal{I}_1 \mathcal{R} u_{j,l} \quad (3.8)$$

$$\mathcal{V}_{j,l}^*(g) := \mathcal{R}^* (v_{j,l}^* * g). \quad (3.9)$$

*Then the system  $(u_{j,l}, \mathcal{V}_{j,l}^*, 2^{-j/2})_{j,l \in \Lambda}$  defines a TI-DFD for  $\mathcal{R}$ , which we will call TI-WVD for the Radon transform.*

*Proof.* For the proof it remains to verify that  $(\mathcal{V}_{j,l}^*)_{j,l}$  satisfies (TI2). Let  $g \in \text{ran}(\mathcal{R})$ , then by the Parseval identity and changing to polar coordinates inside the integral, we have

$$\begin{aligned}
\|\mathcal{V}_{j,l}^* g\|^2 &= \|\mathbf{F}(\mathcal{V}_{j,l}^* g)\|^2 \\
&= \int_{\mathbb{S}^1} \int_0^\infty \frac{1}{\sigma} |\widehat{v}_{j,l}^*(\theta, \sigma)|^2 \cdot |\widehat{g}(\theta, \sigma)|^2 d\sigma d\theta
\end{aligned}$$



$$= \int_{\mathbb{S}^1} \int_0^\infty \frac{\sigma}{2^j} |\widehat{u}_{j,l}(\sigma\theta)|^2 \cdot |\widehat{g}(\theta, \sigma)|^2 d\sigma d\theta.$$

Since  $a2^j \leq \sigma \leq b2^j$  on the support of the integrand and since  $(u_{j,l})_{j,l}$  is a TI frame, taking the sum on both sides yields  $aA \leq \sum_{j,l} \|\mathcal{V}_{j,l}^* g\|^2 \leq bB$ , where  $A, B > 0$  are the TI frame constants of  $(u_{j,l})_{j,l}$ .  $\square$

## 4 Numerical Experiments

In this section, we present numerical comparisons between the classical WVD [4,5] and the TI-WVD for the Radon transform, introduced in this article. The essential step in implementing the filtered TI-DFD reconstruction formula (2.3) is calculating  $\mathcal{V}_{j,l}^* g = \mathcal{R}^*(v_{j,l}^* * g)$ . To this end, we use the explicit expression (3.6) and calculate

$$\mathcal{R}^*(v_{j,l}^* * g) = \kappa_{j,l}(u_{j,l}^* * \mathcal{R}^* \mathcal{I}_1 g).$$

Note that  $\mathcal{R}^* \mathcal{I}_1$  is the filtered backprojection (FBP) as given in (3.5). Together with the Tikhonov filter  $\Phi_\alpha^{(2)}(\kappa) := \kappa/(\kappa^2 + \alpha)$ , the regularized TI-DFD (2.3) for the Radon transform is given as

$$\mathcal{R}_\alpha g = \sum_{j,l} \frac{2^{-j/2}}{2^{-j} + \alpha} u_{j,l} * (u_{j,l}^* * \mathcal{R}^* \mathcal{I}_1 g).$$

Since the wavelet transform and the FBP can be efficiently implemented, this results in an efficient algorithm of the regularized reconstruction. For the implementation, we used Python 3.9.12. The Radon transform and in particular the FBP were implemented via the scikit-image package, version 0.19.2 [16]. The (TI) wavelet transform was employed via the Py-Wavelets package, version 1.3.0 [10].

We use a discretized synthetic phantom  $f \in \mathbb{R}^{256 \times 256}$  and chose the maximum of 8 decomposition levels using the Haar-wavelet as underlying frame. Note that the Haar-wavelet is not band-limited, and thus Theorem 3.5 not applicable in this case. However, we expect the result of the theorem to hold under weaker assumptions, but we do not yet have proof of this. We added white Gaussian noise to the data  $g^\delta = \mathcal{R}f + \delta\eta$ . Here, we chose  $\delta = 0.05$  and  $\eta \sim \|\mathcal{R}f\|_\infty \mathcal{N}(0, 1)$ . To guarantee a fair comparison, we performed a parameter search to determine the optimal regularization parameter  $\alpha > 0$  for both methods. The parameter was optimized in terms of the relative  $\ell^2$  reconstruction error  $\|f_{\text{rec}}^\alpha - f\|_2 / \|f\|_2$ , where  $f$  is the ground truth and  $f_{\text{rec}}^\alpha$  is the reconstruction, depending on the selected parameter.




Figure 4.1: Reconstructions from noisy Radon data  $g^\delta$  with  $\delta = 0.05$ . **(a)** Filtered backprojection reconstruction using (3.5). **(b),(c)** Reconstructions via WVD and TI-WVD, respectively. Both reconstructions use the maximum number of 8 levels of decomposition for the Haar-wavelet.

Numerical results are shown in Figure 4.1, which clearly show that the TI-DFD approach outperforms the standard WVD. While the denoising property of both methods is evident, the decimated DFD suffers from the well-known block like artifacts which are due to the sub sampling step in the decimated wavelet decomposition. Quantitatively, the relative  $\ell^2$  reconstruction error for the WVD amounts to 0.054 and for the TI-WVD to 0.048.

## 5 Conclusion

In this article we presented the concept of the translation invariant frame decomposition (TI-DFD) for the solution of linear operator equations. Subsequently, we constructed a TI wavelet-vagulette decomposition (TI-WVS) for the Radon transform as an instance of the TI-DFD. An advantage of classical frame decompositions is the translation invariance of the system which also has been demonstrated numerically. We have seen, that the use of translation invariant frames leads to improved reconstructions when compared to classical frames.

## Acknowledgments

The contribution by S. G. is part of a project that has received funding from the European Union’s Horizon 2020 research and innovation program under the Marie Skłodowska-Curie grant agreement No 847476. The views and opinions expressed herein do not necessarily reflect those of the European Commission.

## References

- [1] E. J. Candès and D. L. Donoho. Recovering edges in ill-posed inverse problems: optimality of curvelet frames. *Ann. Statist.*, 30(3):784–842, 06 2002.
- [2] R. R. Coifman and D. L. Donoho. *Translation-Invariant De-Noising*, pages 125–150. Springer New York, New York, NY, 1995.
- [3] F. Colonna, G. Easley, K. Guo, and D. Labate. Radon transform inversion using the shearlet representation. *Applied and Computational Harmonic Analysis*, 29(2):232–250, 2010.
- [4] D. L. Donoho. Nonlinear solution of linear inverse problems by wavelet-vaguelette decomposition. *Applied and Computational Harmonic Analysis*, 2(2):101 – 126, 1995.
- [5] A. Ebner, J. Frikel, D. Lorenz, J. Schwab, and M. Haltmeier. Regularization of inverse problems by filtered diagonal frame decomposition, 2020.
- [6] J. Frikel and M. Haltmeier. Sparse regularization of inverse problems by operator-adapted frame thresholding. In *Mathematics of Wave Phenomena*, pages 163–178, Cham, 2020. Springer International Publishing.
- [7] S. Göppel, J. Frikel, and M. Haltmeier. Translation invariant diagonal frame decomposition of inverse problems and their regularization, 2022.
- [8] A. N. H. Engl, M. Hanke. *Regularization of inverse problems*. Mathematics and Its Applications. Springer, 1 edition, 1996.
- [9] S. Hubmer, R. Ramlau, and L. Weissinger. On regularization via frame decompositions with applications in tomography. *Inverse Problems*, 38(5):055003, 2022.
- [10] G. R. Lee, R. Gommers, F. Waselewski, K. Wohlfahrt, and A. O’Leary. Pywavelets: A Python package for wavelet analysis. *Journal of Open Source Software*, 4(36):1237, 2019.
- [11] S. Mallat. *A Wavelet Tour of Signal Processing, Third Edition: The Sparse Way*. Academic Press, Inc., USA, 3rd edition, 2008.
- [12] G. P. Nason and B. W. Silverman. *The Stationary Wavelet Transform and some Statistical Applications*, pages 281–299. Springer New York, New York, NY, 1995.
- [13] F. Natterer. *The Mathematics of Computerized Tomography*. Classics in Applied Mathematics. Society for Industrial and Applied Mathematics, 2001.

- [14] K. T. Smith, D. C. Solmon, and S. L. Wagner. Practical and mathematical aspects of the problem of reconstructing objects from radiographs. *Bulletin of the American Mathematical Society*, 83(6):1227–1270, 1977.
- [15] D. C. Solmon. The x-ray transform. *Journal of Mathematical Analysis and Applications*, 56(1):61–83, 1976.
- [16] S. van der Walt, J. L. Schönberger, J. Nunez-Iglesias, F. Boulogne, J. D. Warner, N. Yager, E. Gouillart, T. Yu, and the scikit-image contributors. scikit-image: image processing in Python. *PeerJ*, 2:e453, 6 2014.